

A Brief Introduction to Finite Element Methods for Fluid Flow Problems

James Percival

AMCG, Department of Earth Science and Engineering Imperial College London

Fluidity Training 2014

글 🕨 🖌 글 🕨

AMCG

Outline

Numerical Solution of PDEs

Standard numerical methods for PDEs.

Finite Elements from First Principles

Example: Poisson equation Strong and Weak Forms Boundary Conditions

Extending the method

Two and Three dimensional FEM High order polynomials and discontinous FEM

Spaces, Forms and Functions

More on Vector Spaces Existence, Uniqueness and Convergence of Solutions

A Note on Implementation



→ < ∃→

Numerical Solutions to PDEs

Mathematical descriptions of the equations governing physical processes are often in the form of partial differential equations (PDEs). Variable values are functions of time and satisfy relationships between variables and their partial derivatives.

Examples

$$\frac{\partial T}{\partial t} = \kappa \frac{\partial^2 T}{\partial x^2}, \qquad (\text{heat equation})$$
$$\frac{\partial^2 a}{\partial t^2} = c^2 \frac{\partial^2 a}{\partial x^2} \qquad (\text{wave equation})$$
$$\frac{\partial \tau}{\partial t} + u \frac{\partial \tau}{\partial x} = \kappa \frac{\partial^2 \tau}{\partial^2 x} \qquad (\text{advection-diffusion equation})$$

イロト 不得下 不同下 不同下

AMCG

Numerical Solutions to PDEs

Solving PDEs assumes knowledge of the variables at an infinite number of locations in space and time. However memory available in computers (and humans) is finite. Equations must be discretized to create a smaller (and easier to solve) problem. Eg. to a matrix problem like

$$\begin{pmatrix} 3 & 2 & 0 \\ 4 & 4 & 1 \\ 0 & 2 & 3 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 1 \\ 3 \\ 2 \end{pmatrix},$$

or, in a more general form,

$$Ax = b.$$

AMCG

Numerical Solutions to PDEs

Some standard techniques for discretization are:

- 1. Finite difference methods.
- 2. Finite volume methods.
- 3. Spectral methods.
- 4. Finite element methods.

The Fluidity model primarily implements the last of these.





Finite Difference Methods

- Reduce problem domain to finite set of points.
- Replace exact derivatives with approximate difference equations.



x

$$\frac{df}{dx} \approx \frac{\Delta f}{\Delta x} = \frac{f(x_{i+1}) - f(x_{i-1})}{x_{i+1} - x_{i-1}}$$

▲□▶▲圖▶▲圖▶▲圖▶ ▲□▶



Finite Volume Methods

- Break problem domain into a finite set of sub-\emph{volumes}.
- Solve for volume integral of quantities inside. Typically reduces problem into flux calculation across faces of the volume.
- If fluxes depend on derivatives then another method (e.g. finite differences) must be used to find them.

$$\frac{d}{dt} \int_{\Omega_i} \rho dV = \sum_{\text{faces}} \int_{\delta \Omega_i^{(j)}} \rho \boldsymbol{u} \cdot \boldsymbol{n} dS$$



MCG

Spectral Methods

- Represent variables as (limit of) sum of orthogonal basis functions.
- Global basis functions vary over entire domain.
- Truncate infinite series and calculate behaviour of finite set of coefficients.

$$f(x) = \sum_{n=1}^{\infty} a_n \cos\left(n\frac{\pi x}{L}\right) + b_n \sin\left(n\frac{\pi x}{L}\right)$$
$$\approx \sum_{n=1}^{N} a_n \cos\left(n\frac{\pi x}{L}\right) + b_n \sin\left(n\frac{\pi x}{L}\right)$$





Spectral Methods



‼= ୬९୯ ∰€**⊊**



Spectral Methods



‼= ୬९୯ ∰€**⊊**



Spectral Methods



‼= ୬९୯ ∰€**⊊**

Spectral Methods



‼= ୬९୯ ∰€**⊊**



Finite Element Methods

 Break the problem domain down into a finite set of sub-volumes (elements).

AMCG

- Represent variables as a sum of basis functions.
- Basis functions non-zero only on local set of subvolumes.
- ► Solve integral equation form of PDE.

Hybrid Methods

Very common to combine these different approaches

- Couple finite volume method for globally conserved quanity to finite difference method to calculate fluxes
- Couple finite volume method for globally conserved quanity to finite element method to calculate fluxes. "Control Volume" method
- Fluidity velocity solver: Finite element method in space, finite difference in time

글 🖌 🖌 글 🕨

MCG

Poisson Equation: Pressure in Navier-Stokes

Incompressible Navier-Stokes equations

$$\frac{\partial u}{\partial t} + u \cdot \nabla u = -\nabla p + \nu \nabla^2 u, \qquad (\text{momentum})$$
$$\nabla \cdot u = 0, \qquad (\text{continuity})$$

Taking divergence

$$\underbrace{\frac{\partial}{\partial t} \left(\nabla \cdot \boldsymbol{u} \right)}_{=0} + \nabla \cdot \left(\boldsymbol{u} \cdot \nabla \boldsymbol{u} \right) = -\nabla^2 p + \underbrace{\nu \nabla^2 \nabla \cdot \boldsymbol{u}}_{=0}.$$

I.e.

$$\nabla^2 p = -\nabla \cdot (\boldsymbol{u} \cdot \nabla \boldsymbol{u}) \,.$$



글 🕨 🖌 글 🕨

Poisson Equation

General form of this equation

Poisson Equation

$$\nabla^{2}\psi+f(\boldsymbol{x})=0,\quad\forall\boldsymbol{x}\in\Omega$$

In 1D, setting Ω to the unit interval:

$$\frac{\partial^2 \psi}{\partial x^2} + f(x) = 0 \quad \forall x \in (0, 1).$$

글 🕨 🖌 글 🕨

AMCG

This is the strong form of the Poisson equation.

Strong Form vs. Weak Form of an Equation

Strong form

Equation (*) true individually for each point in space,

 $\nabla^2 \psi + f(x) = 0,$

for all x in domain Ω . Test a ψ by checking equation holds individually for each point in space.

Weak form

Integral equation holds for all choices of a 'test function', ϕ ,

$$\int_{\Omega} \phi\left(\nabla^2 \psi + f\right) \, dV = 0,$$

where $\phi: \Omega \to \mathbb{R}$ is from a function space defined later. Test possible ψ ('trial function') by checking integral equation holds for all test functions, ϕ .



Quick Review of Vector Spaces

A set, $\mathcal V,$ is a vector space if it has + (addition) and scalar multiplication operators where

$$a + (b + c) = (a + b) + c$$
, (associativity)
 $a + b = b + a$, (commutivity)

there exists $\mathbf{0} \in \mathcal{V}$ such that $a + \mathbf{0} = a$, for all $a \in \mathcal{V}$,

for all a there exists $-a \in \mathcal{V}$ such that $a + (-a) = \mathbf{0}$,

$$\begin{aligned} &\alpha \left(a+b \right) = \alpha a + \alpha b, & (\text{distributivity I}) \\ &\left(\alpha + \beta \right) a = \alpha a + \beta a, & (\text{distributivity II}) \\ &\alpha \left(\beta a \right) = \left(\alpha \beta \right) a, \\ &1a = a. \end{aligned}$$

글 🖌 🔺 글 🕨

AMCG

Functions are a vector space under the following definitions of addition and scalar multiplication:

$$(f+g)(x) = f(x) + g(x),$$

(αf)(x) = α ($f(x)$).

문▶ ★ 문▶

AMCG

I.e. functions are added/multiplied pointwise based on their result. Necessary axioms all follow from the normal rules of addition/multiplication.





≣I≡ ୬۹୯ MCC





EIE 996





≣ા= ୬**૧**୯ ∰દ**િ**∼



Within the space of functions there are many smaller subspaces. Eg:

AMCG

Examples

Polynomials $f(x) = 1 + 3x + 4x^2 + 5x^3$ Functions on an interval $f(x) = \begin{cases} 0 & x < 0, \\ e^x & 0 \le x \le 1, \\ 0 & x > 1. \end{cases}$ Twice differentiable functions $f(x) = \begin{cases} x^2 + 2, & x < 0, \\ 2(e^x - x), & x \ge 0. \end{cases}$

Vector Spaces

Key idea of finite element method:

- Desire an exact solution ψ, from an inifinite dimensional vector space, V, which satisfies weak form equation, for test functions φ in U.
- Find ψ^δ in approximate vector (sub)space V^δ ⊂ V with finite representation, which satisfies same weak form equation, for test functions φ^δ in approximate space U^δ ⊂ U.

글 🖌 🖌 글 🕨

MCG

• Expectation that as $\delta \to 0$, $\mathcal{V}^{\delta} \to \mathcal{V}$ and $\psi^{\delta} \to \psi$.

Strong and Weak Forms

A solution to the strong form of the equations will be a solution to the weak form equations: If $\nabla^2\psi+f=0$ then

$$\int \phi \left(\nabla^2 \psi + f \right) \, dV = \int \phi \cdot 0 \, dV = 0,$$

independent of ϕ , i.e. for any possible choice of test space.



글 🖌 🖌 글 🕨

Strong and Weak Forms

A solution to the weak form of the equations may be a solution to the strong equations if it is smooth enough.

The weak formulation extends the equations to allow non-smooth solutions which exist in a distributional sense.

Examples of common distributions

$$\delta(x), \quad \int_{-\infty}^{a} f(x) \,\delta(x) = \begin{cases} f(0), & a > 0, \\ 0, & a < 0. \end{cases}$$
(Dirac delta)
$$H(x) := \int_{-\infty}^{x} \delta(s) \, ds = \begin{cases} 0, & x < 0, \\ 1, & x > 0. \end{cases}$$
(Heaviside)

AMCG

Example of a Weak Nonclassical Solution

$$\psi - \nabla^2 \psi (x) = a\delta (X - x)$$
(Helmholtz)
$$\psi = \begin{cases} a \exp (x - X) & x \le X, \\ a \exp (X - x) & x > X \end{cases}$$







Review of Section

- Strong form of PDEs prescribes behaviour pointwise
 - Finite difference methods work at a finite number of points
- Weak form of PDEs prescribes behaviour over intervals (areas, volumes etc.)
 - Finite element methods work over a finite number of intervals
- ► Functions live in vector spaces, which can be approximated.

Boundary Conditions for Weak Equations

Two possible forms of boundary condition for a solution to the Poisson equation to be well posed:

1. Dirichlet:
$$\psi(x) = a(x)$$
 for $x \in A \subset \delta\Omega$,

2. Neumann:
$$\frac{\partial \psi}{\partial x} = b(x)$$
 for $x \in B \subset \delta \Omega$.

In the Galerkin formulation, Dirichlet boundary conditions typically require explicit modification of the structure of the problem to be solved, whereas Neumann conditions are dealt with naturally as part of the formulation. We'll use a Dirichlet boundary condition at x = 0, and a Neumann condition at x = 1 in our example.



글 🖌 🖌 글 🕨

Natural Boundary Conditions

Our weak form equation is

$$\int_0^1 \phi\left(\frac{\partial^2 \psi}{\partial x^2} + f\right) \, dx = 0,$$

Integrate by parts,

$$\int_0^1 \frac{\partial \phi}{\partial x} \frac{\partial \psi}{\partial x} \, dx - \int_0^1 \phi f \, dx = -\left[\phi \frac{\partial \psi}{\partial x}\right]_0^1$$

Chose ϕ to vanish on Dirichlet boundaries (and set $\psi = a(0)$), and use our knowlege of $\frac{\partial \psi}{\partial x}$ on Neumann boundaries:

$$\int_{0}^{1} \frac{\partial \phi}{\partial x} \frac{\partial \psi}{\partial x} \, dx = \int_{0}^{1} \phi f \, dx - \phi \left(1\right) b \left(1\right)$$



▲ 문 ▶ ▲ 문 ▶ _ 문 범

Dirichlet Boundary Conditions

Dirichlet boundary conditions can be enforced through splitting solution into two parts

 $\psi = \psi_0 + \psi_d$

where ψ_d is any (chosen) function satisfying

$$\psi_d(0) = a(0), \qquad \frac{\partial \psi_d}{\partial x}(1) = 0$$

while ψ_0 satisfies a modified weak equation, with ψ_d on r.h.s,

$$\psi_0(0)=0,$$

$$\int_{0}^{1} \frac{\partial \phi}{\partial x} \frac{\partial \psi_{0}}{\partial x} dx = \int_{0}^{1} \phi f \, dx - \phi \left(1\right) b \left(1\right) - \int_{0}^{1} \frac{\partial \phi}{\partial x} \frac{\partial \psi_{d}}{\partial x} dx,$$

Note ψ_0 vanishes on the Dirichlet boundaries, same condition we apply to ϕ .



Boundary Conditions: Strong vs. Weak

More generally, implementations of finite element boundary conditions come in two flavours, strong and weak.

Strong form bc.s

Information contained in boundary condition appears implicitly in the weak form of the PDE. Solve by lifting method.

Weak form bcs.

Information contained in boundary condition appears explicitly in surface integrals in the weak form of the PDE. Solve by direct substitution.

듣▶ ★ 돋▶

Boundary Conditions: Strong vs. Weak

Which method to apply depends on the details of both the original PDE and the weak form to be solved. Sometimes both are possible. For example, consider solving the advection equation,

$$\frac{\partial \tau}{\partial t} + \boldsymbol{a} \cdot \nabla \tau = 0,$$

for a tracer τ , given a known velocity field, a and dirichlet boundary conditions at a inlet.

Note the odd number of spatial derivatives, whereas even number for Poisson equation.

글 🖌 🔺 글 🛌

Boundary Conditions: Strong vs. Weak

Solve

$$\int_{\Omega} \phi\left(\frac{\partial \tau}{\partial t} + \boldsymbol{a} \cdot \nabla \tau\right) dx = 0,$$

Dirichlet bcs must be applied strongly.

Weak form bcs.

Integrate by parts,

$$\int_{\Omega} \phi \frac{\partial \tau}{\partial t} - \tau \nabla \cdot (\phi a) \, dx$$
$$= -\int_{\delta \Omega^+} a \cdot n \phi \tau_b \, dS,$$

where τ_b is the Dirichlet bc, applied weakly.

E ► ★ E ► .

When using this sort of weak boundary condition, values may not be quite what you'd expect, however fluxes should be right.

Finite Element Basis Functions

Need discrete finite dimensional representation of problem to do numerical calculations on a computer. Set

$$\psi^{\delta}(x) = \sum_{i=1}^{N} \hat{\psi}_{i} N_{i}(x)$$

where $\hat{\psi}_i \in \mathbb{R}$ is a scalar parameter and $N_i : \Omega \to \mathbb{R}$ is a fixed shape function specifing spatial dependence. Can do the same for the space of test functions, ϕ .



글 🖌 🖌 글 🕨

Finite Element Functions

For our 1D Poisson equation example we can choose to use the set of continuous, piecewise linear functions ('shape functions') on subdivisions of the unit interval.

$$N_i = egin{cases} 0, & x \leq x_{(i-1)}, \ rac{x-x_{(i-1)}}{x_i-x_{(i-1)}}, & x_{(i-1)} < x \leq x_i, \ rac{x_{(i+1)}-x}{x_{(i+1)}-x_i}, & x_i < x \leq x_{(i+1)}, \ 0. & x > x_{(i+1)}. \end{cases}$$

Functions are equal to 1 at the set of points $[0, x_1, x_2, \dots, x_{n-1}, 1]$, sometimes called 'nodes' or 'degrees of freedom'. The subdivisions of Ω over which the N_i s are smooth are often called elements.






≣ = ୬.୯ ∰©



ے اے ان مرد میں کو میں



APC ≓IE MCC



ন্দ লাভ নি দিন্দ্র



≣ા= ୬**૧**୯ ∰દ**≪**





SPC ≓IE MCC





≣ा= ୬९୯ ∰¢द≪

Galerkin Approximation

To obtain the Galerkin approximation of the Poisson equation, we find the (unique) solution of the weak form equation when ψ and ϕ are approximated by our finite element expansions,

$$egin{aligned} \psi^\delta &= \sum_{i=0}^n \hat{\psi}_i N_i, \ \phi^\delta &= \sum_{j=0}^n \hat{\phi}_j N_j. \end{aligned}$$

The ψ^{δ} are called trial functions and the function space they come from is the trial space. The ϕ^{δ} are called test functions and live in the trial space. Computation involves obtaining the finite number of $\hat{\psi}_i$. Can then be solved on a computer.

Galerkin Approximation

Substituing the finite representations into (*) we get

$$\int \sum_{j=1}^{n} \hat{\phi}_{j} \frac{\partial N_{j}}{\partial x} \sum_{i=1}^{N} \hat{\psi}_{i} \frac{\partial N_{i}}{\partial x} dV + v_{N}^{\delta} b(x_{N}) = \int \sum_{j=1}^{n} \hat{\phi}_{j}^{\delta} N_{j} f dV,$$
$$\hat{b}_{j} \left(\underbrace{\left[\int \frac{\partial N_{j}}{\partial x} \frac{\partial N_{i}}{\partial x} dV \right]}_{\text{matrix } D_{ij}} \hat{\psi}_{i} - \int f N_{j} dV + \begin{cases} 0, & j = 1, \dots, n-1 \\ b(x_{n}), & j = n \end{cases} \right)$$
$$= 0$$

If bracket vanishes, solution applies for any $\hat{\phi}_j$, so we can drop them.

The Right Hand Side

Generally the right hand side of the equation is known explicitly as a function $f: \Omega \to \mathbb{R}$. Hence $\int_0^1 \phi^{\delta} f \, dx$ can be calculated exactly. In practice (especially for coupled problems,) it is usually represented in the approximate function space,

$$f^{\delta}\left(x
ight)=\sum_{i=1}^{N}\widehat{f}_{i}N_{i}\left(x
ight)$$
 ,

where, for our choice of shape functions,

$$\hat{f}_i = f(x_i).$$

Imperial College London

Finite Element Poisson Matrix Problem Dirichlet condition: $\hat{\psi}_0 = a(0)$, turns up on right hand side:

$-\frac{2}{h}$	$\frac{1}{h}$	0	0	0	0	ψ_1		$\frac{\frac{h}{6}\hat{f}_0 + \frac{2h}{3}\hat{f}_1 + \frac{h}{6}\hat{f}_2 - \frac{1}{h}a(0)$
$\frac{1}{h}$	$-\frac{2}{h}$	$\frac{1}{h}$	0	0	0	ψ_2		$\frac{h}{6}\hat{f}_1 + \frac{2h}{3}\hat{f}_2 + \frac{h}{6}\hat{f}_3$
0	$\frac{1}{h}$	$-\frac{2}{h}$	$\frac{1}{h}$	0	0	ψ_3		$\frac{h}{6}\hat{f}_2 + \frac{2h}{3}\hat{f}_3 + \frac{h}{6}\hat{f}_4$
0	0	$\frac{1}{h}$	$-\frac{2}{h}$	$\frac{1}{h}$	0	ψ_4	_	$\frac{h}{6}\hat{f}_3 + \frac{2h}{3}\hat{f}_4 + \frac{h}{6}\hat{f}_5$
0	0	0	$\frac{1}{h}$	$-\frac{2}{h}$	$\frac{1}{h}$	ψ_5		$\frac{h}{6}\hat{f}_4 + \frac{2h}{3}\hat{f}_5 + \frac{h}{6}\hat{f}_6$
0	0	0	0	$\frac{1}{h}$	$-\frac{1}{h}$	ψ_6		$rac{h}{6}\hat{f}_5+rac{h}{3}\hat{f}_6$ -b(1)

Local to global

Note that

$$egin{aligned} D_{ij} &= \int_\Omega rac{\partial N_i}{\partial x} rac{\partial N_j}{\partial x} \, dV, \ &= \sum_k \int_{\Omega^{(k)}} rac{\partial N_i}{\partial x} rac{\partial N_i}{\partial x} \, dV, \end{aligned}$$

where the $\Omega^k = [x_i, x_{i+1}]$ only contribute if

$$\frac{\partial N_i}{\partial x} \neq 0, \quad \frac{\partial N_j}{\partial x} \neq 0.$$

▶ < 글 ▶ .

AMCG

Global matrix assembled from sum of local integrals over elements.

Finite Difference Poisson Matrix Problem

Finite Difference discretization of same problem:

Finite Volume Poisson Matrix Problem

Finite volume discretization of same problem. 2 point finite difference approximation for flux terms:

$\frac{-3}{h}$	$\frac{1}{h}$	0	0	0	0	$\psi_{\frac{1}{2}}$	$\begin{array}{c} \psi_{\frac{1}{2}} \\ \psi_{\frac{3}{2}} \\ \psi_{\frac{5}{2}} \\ \psi_{\frac{7}{2}} \\ \psi_{\frac{7}{2}} \\ \psi_{\frac{9}{2}} \end{array} =$	$h\hat{f}_{1/2} - 2\frac{a(0)}{h}$
$\frac{1}{h}$	$\frac{-2}{h}$	$\frac{1}{h}$	0	0	0	$\psi_{\frac{3}{2}}$		$h\hat{f}_{3/2}$
0	$\frac{1}{h}$	$\frac{-2}{h}$	$\frac{1}{h}$	0	0	$\psi_{\frac{5}{2}}$		$h\hat{f}_{5/2}$
0	0	$\frac{1}{h}$	$\frac{-2}{h}$	$\frac{1}{h}$	0	$\psi_{\frac{7}{2}}$		$h \hat{f}_{7/2}$
0	0	0	$\frac{1}{h}$	$\frac{-2}{h}$	$\frac{1}{h}$	$\psi_{\frac{9}{2}}$		$h\hat{f}_{9/2}$
0	0	0	0	$\frac{1}{h}$	$\frac{-1}{h}$	$\psi_{\frac{11}{2}}$		$h\hat{f}_{11/2}-b(1)$

Solutions: Finite Element $f = 10 \sin(5x) + \frac{1}{2} \cos(3(x + \frac{1}{2})), \psi(0) = 0, \frac{d\psi}{dx} = 1:$



Intro to FEM for Fluids; James Percival

Solutions: Finite Difference $f = 10 \sin(5x) + \frac{1}{2} \cos(3(x + \frac{1}{2})), \psi(0) = 0, \frac{d\psi}{dx} = 1:$



Solutions: Finite Volume $f = 10 \sin(5x) + \frac{1}{2} \cos(3(x + \frac{1}{2})), \psi(0) = 0, \frac{d\psi}{dx} = 1:$



Example II: Advection-Diffusion

Use same finite element framework for 1D advection diffusion equation:

$$\frac{\partial \tau}{\partial t} + \frac{\partial}{\partial x} \left(u\tau \right) = \frac{\partial}{\partial x} \left(\kappa \frac{\partial \tau}{\partial x} \right)$$
$$\int_0^1 \phi \frac{\partial \tau}{\partial t} \, dx = \int \frac{\partial \phi}{\partial x} \left(u\tau - \kappa \frac{\partial \tau}{\partial x} \right) \, dx$$

E ► ★ E ► .

AMCG

Example II: Advection-Diffusion

Given FEM framework, crank the handle to reduce the problem; Discretize through Finite Element Galerkin Method,

$$egin{aligned} & au^{\delta} = \sum_{i=0}^n \hat{ au}_i N_i^{ au}, \ & u^{\delta} = \sum_{i=0}^n \hat{u}_i N_i^u \ & \phi^{\delta} = \sum_{i=0}^n \hat{\phi}_i N_i^{ au}, \end{aligned}$$

Note that the method doesn't require $N_i^u = N_i^{\tau}$. Mixed formulations are possible

★ 문 ▶ ★ 문 ▶ _ 문

Example II: Advection-Diffusion

Following substitution, integrate by parts to obtain

$$\underbrace{\int_{0}^{1} N_{i} N_{j} dx}_{\text{Mass matrix}^{"} M_{ij}} \frac{\partial \hat{\tau}_{j}}{\partial t} - \int_{0}^{1} \frac{\partial N_{i}}{\partial x} \left(N_{j} \sum_{k=0}^{n} u_{k} N_{k}^{u} - \kappa \frac{\partial N_{j}}{\partial x} \right) dx \hat{\tau}_{j} = 0,$$

or in matrix form,

$$\mathbf{M}\frac{\partial \boldsymbol{\tau}}{\partial t} + \mathbf{A}\left(\boldsymbol{u}\right)\boldsymbol{\tau} + \mathbf{D}\left(\boldsymbol{\kappa}\right)\boldsymbol{\tau} = \boldsymbol{0},$$

AMCG

This is the FEM form of the tracer advection-diffusion equation. Further details will depend on the choice of shape functions and timestepping method.



Review of Section

- Finite element methods solve weak (integral) equations
- Functions get approximated by finite dimensional summations of functions, non-zero over small regions of problem domain (elements)
- Linear PDE problem gives a linear (matrix) problem for the $\hat{\psi}_i$.
- The efficient computational representation and solution of these sorts of problems will form the basis of the other session.

→ < Ξ→















To increase the number of free parameters in the approximate solution space (and thus attempt to get a more accurate solution) there are several options:

- ▶ More, smaller subdivisions [step size,*h*]
 - This is the system used in Fluidity's mesh adaptivity.
- Use higher order polynomials, e.g. quadratic functions [polynomial order, p]
- Use discontinuous functions [Discontinuous Galerkin formulation]

듣▶ ★ 돋▶ .

- ▶ Projection (in black) of smooth function (red).
- ► Linear, continuous basis, Galerkin method (P1 CG).
- \blacktriangleright In 1d degrees of freedom \approx no. of elements



- ▶ Projection (in black) of smooth function (red), & error (blue).
- ► Linear, continuous basis, Galerkin method (P1 CG).
- \blacktriangleright In 1d degrees of freedom \approx no. of elements



- May increase the number of elements
- More elements mean more degrees of freedom
- One of the methods used in Fluidity adaptivity routines



Quadratic shape functions

$$N_{2i} = \begin{cases} 0, & x \leq x_{i-1}, \\ \frac{2x^2 - (3x_{i-1} + x_i)x + (x_i + x_{i-1})x_{i-1}}{(x_i - x_{i-1})^2}, & x_{i-1} < x \leq x_i, \\ \frac{2x^2 - (3x_{i+1} + x_i)x + (x_i + x_{i+1})x_{i+1}}{(x_{i+1} - x_i)^2}, & x_i < x \leq x_{i+1}, \\ 0, & x > x_{i+1}. \end{cases}$$

$$N_{2i+1} = \begin{cases} 0, & x < x_i, \\ -\frac{x^2 - (x_{i+1} + x_i)x + x_ix_{i+1}}{(x_{i+1} - x_i)^2}, & x_i < x \leq x_{i+1}, \\ 0, & x > x_{i+1}. \end{cases}$$



- ▲ 문 ► - ▲ 문 ►





≣⇒

AMCG





∃⇒

AMCG





≣⇒

AMCG





≣⇒

AMCG





∃⇒

AMCG

- Projection of a smooth function.
- ► Quadratic, continuous basis, Galerkin method (P2 CG).
- \blacktriangleright In 1d degrees of freedom \approx 2 \times no. of elements.
- Good representation of slowly varying functions


Increasing the degrees of freedom

Discontinuous linear shape functions

$$N_{2i} = \begin{cases} 0, & x \leq x_i, \\ \frac{(x_{i+1}-x)}{(x_{i+1}-x_i)}, & x_i < x \leq x_{i+1}, \\ 0. & x > x_{i+1}. \end{cases}$$

$$N_{2i+1} = \begin{cases} 0, & x < x_i, \\ -\frac{(x-x_i)}{(x_{i+1}-x_i)}, & x_i < x \le x_{i+1}, \\ 0, & x > x_{i+1}. \end{cases}$$

★ E ► ★ E ►

AMCG

< 口 > < 同

Increasing the degrees of freedom

- Projection of a smooth function.
- ► Linear, discontinuous basis, Galerkin method (P1 DG).
- \blacktriangleright In 1d degrees of freedom $\approx\!\!2\times$ no. of elements.
- Good representation of discontinuties/fronts/large gradients.





Review of Section

- The degrees of freedom (and thus size) of a problem can be increased by:
 - increasing the number of element subdivisions (making step size h smaller)
 - increasing the order of the shape functions applied on elements (increasing polynomial degree, p)
 - relaxing continuity constraints at the interface between elements (discontinuous Galerkin method, nonconforming elements)

★ 문 ► ★ 문 ► .

MCG



Reassuring Mathematics

This section summarises some useful results from mathematical analysis for finite element problems. In particular, we note that results exists to show that, under certain provisos finite element solutions to a given problem

- exist
- are unique
- converge
- converge to the right answer.



Choice of Vector Spaces

Going back to the weak form for the original infinite dimensional problem,

$$\int_{\Omega} \nabla \phi \cdot \nabla \psi \, dV = \int_{\Omega} \phi f \, dV + \int_{\delta \Omega^N} \phi b \, dV,$$

글 🖌 🔺 글 🛌

AMCG

it is obvious that ψ and $\nabla\psi$ must be well behaved enough for these integrals to exist.

Choice of Vector Spaces

We require the function is square integrable,

$$\|\psi\|^2 := \int_{\Omega} \psi^2 \, dV < \infty. \tag{1}$$

(The space of function which satisfy this is normally called $\mathcal{L}_{2}\left(\Omega\right)$) and also that

$$\|\nabla\psi\|^{2} := \int_{\Omega} \nabla\psi \cdot \nabla\psi \, dV < \infty, \tag{2}$$

글 🖌 🔺 글 🕨

MCG

Functions which satisfy both (1) & (2) are in the space of square integrable functions with square integrable derivatives, denoted $\mathcal{H}^1(\Omega)$. This is a Sobolev space.

Weak equations and Bilinear forms

The volume integral in (*) defines a symmetric bilinear form, $a: \mathcal{H}^{1}(\Omega) \times \mathcal{H}^{1}(\Omega) \rightarrow \mathbb{R}$,

$$a(\phi,\psi):=\int_{\Omega}\nabla\psi\cdot\nabla\phi\,d^nx.$$

where

$$a(\phi, \psi) = a(\psi, \phi),$$

$$a(c_1\phi + c_2\xi, \psi) = c_1a(\phi, \psi) + c_2a(\xi, \psi).$$

글 🖌 🖌 글 🕨



Lax-Milgram Theorem

Two properties of the bilinear form, a, are used to show well-posedness:

$$\begin{split} a(\phi,\psi) &\leq C \|\psi\| \|\phi\| \text{ for some C} > 0, \qquad \text{(continuity)} \\ a(\psi,\psi) &\geq c \|\psi\|^2 \text{ for some c} > 0. \qquad \text{(coercive/elliptic)} \end{split}$$

글 🖌 🖌 글 🕨

Well posedness - existence

Existence follows from application of Riesz representation theorem to the Hilbert space problem

$$a\left(u,v
ight) =f\left(v
ight)$$
 ,

Precise details lie outside of the scope of this lecture, but effectively guarantees an "inverse" to the map

$$\phi_{v}\left(u\right)=a\left(u,v\right),$$

ㅋㅋ ㅋㅋㅋ

so that for sufficiently smooth data we can always get a solution.

Well posedness - uniqueness:

Suppose there are two different solutions, ψ_1 and ψ_2 , i.e.

$$a\left(\phi,\psi_{1}\right)=a\left(\phi,\psi_{2}\right)=\int_{\Omega}\phi f\,d^{n}x,\text{ for all }\phi\in\mathcal{H}^{1}\left(\Omega
ight).$$

Then

$$a\left(\phi,\psi_1-\psi_2\right)=0$$

but $\psi_{1}-\psi_{2}\in\mathcal{H}^{1}\left(\Omega
ight)$, so can choose to test $\phi=\psi_{1}-\psi_{2}$ Then

$$a(\psi_1 - \psi_2, \psi_1 - \psi_2) = 0 \ge c \|\psi_1 - \psi_2\|^2$$
,

So $\psi_1 = \psi_2$, hence solution is unique.



(《문》 《문》 문)님

Well posedness - convergence

Let $\psi \in \mathcal{V}$ be exact solution, $\psi^{\delta} \in \mathcal{V}^{\delta} \subset \mathcal{V}$ be the finite element solution $\xi \in \mathcal{V}^{\delta}$ be an arbitrary function. Then $\psi^{\delta} - \xi \in \mathcal{V}$ and $\psi^{\delta} - \xi \in \mathcal{V}^{\delta}$ and

$$a(\underbrace{\psi^{\delta} - \xi}_{\in \mathcal{V}}, \psi) = \int_{\Omega} \left(\psi^{\delta} - \xi\right) f d^{n} x \qquad (\text{From PDE})$$

$$a(\underbrace{\psi^{\delta}-\xi}_{\in\mathcal{V}^{\circ}},\psi^{\delta}) = \int_{\Omega} \left(\psi^{\delta}-\xi\right) f d^{n}x$$
 (FEM)

글 🕨 🖌 글 🕨



Well posedness - convergence

$$\begin{split} c \left\| \psi - \psi^{\delta} \right\|^{2} &\leq a \left(\psi - \psi^{\delta}, \psi - \psi^{\delta} \right), \\ &= a \left(\psi - \psi^{\delta}, \psi - \psi^{\delta} \right) + a (\psi - \psi^{\delta}, \psi^{\delta} - \xi) \\ &+ a \left(\psi^{\delta} - \xi, \psi^{\delta} \right) - a \left(\psi^{\delta} - \xi, \psi \right), \\ &= a \left(\psi - \psi^{\delta}, \psi - \psi^{\delta} + \psi^{\delta} - \xi \right) \\ &- \int_{\Omega} \left(\psi^{\delta} - \xi \right) f d^{n} x + \int_{\Omega} \left(\psi^{\delta} - \xi \right) f d^{n} x, \\ &= a \left(\psi - \psi^{\delta}, \psi - \xi \right) \leq C \left\| \psi - \psi^{\delta} \right\| \left\| \psi - \xi \right\|. \end{split}$$

Well posedness - convergence

Hence it is guaranteed (Cea's lemma)

$$\left\| \psi - \psi^{\delta}
ight\| \leq rac{C}{c} \inf_{\xi \in \mathcal{V}^{\delta}} \left\| \psi - \xi
ight\|.$$

Choose ξ to be linear projection of ψ , i.e $\xi(x_i) = \psi(x_i)$, $\frac{\partial^2 \xi}{\partial x^2} = 0$, then

$$\|\psi - P\psi\| \le \alpha \sup_{\Omega_i} h_i \sup_{x \in \Omega} \left| \frac{\partial^2 \psi}{\partial x^2} \right|$$

Hence

$$ig|\psi-\psi^{\delta}ig\|\leq rac{lpha C}{c} \sup_{\Omega_{i}} h_{i} \sup_{x\in\Omega} ig|rac{\partial^{2}\psi}{\partial x^{2}}ig| \ ig\|\psi-\psi^{\delta}ig\| o 0 ext{ as } \sup_{\Omega_{i}} h_{i} o 0.$$



→ ▲ 글 ▶



Review of Section

We have shown that finite element approximations to the solutions to $\ensuremath{\mathsf{PDEs}}$

- are unique,
- converge,
- and converge to the right answer.

We have also given a hint that they exist. We have also shown that by using knowledge about the form of the solution we can choose elements to minimize the estimated error for a given number of degrees of freedom.



→ < Ξ→

Numerical method to calculate/approximate integrals:

$$\int_{a}^{b} f(\mathbf{x}) d^{n} \mathbf{x} \approx \sum_{x=1}^{N} w_{i} f(\mathbf{p}_{i})$$



Numerical method to calculate/approximate integrals:

$$\int_{a}^{b} f(\mathbf{x}) d^{n} \mathbf{x} \approx \sum_{x=1}^{N} w_{i} f(\mathbf{p}_{i})$$





Some famous quadratures:

1. Midpoint rule [one point method]

$$w=a-b$$
, $p=rac{a+b}{2}$,

2. Simpson's rule [3 point method]

$$w_1 = \frac{a-b}{6}, \quad w_2 = \frac{4(a-b)}{6}, \quad w_3 = \frac{a-b}{6},$$

 $p_1 = a, \quad p_2 = \frac{a+b}{2}, \quad p_3 = b,$



ㅋㅋ ㅋㅋㅋ

A quadrature is called exact for functions where

$$\int_{a}^{b} f(\mathbf{x}) d^{n} \mathbf{x} = \sum_{x=1}^{N} w_{i} f(\mathbf{p}_{i}).$$

The degree (or order) of a quadrature rule over an interval is the largest integer, n such that method is exact for all

$$p_n = a_0 x^n + a_1 x^{n-1} + \dots a_n \quad a_i \in \mathbb{R}.$$

For a FE method, would like a quadrature which captures the "worst" order of the terms in the integral, eg

$$\int_{x_j}^{x_{j+1}} \nabla N_i^{(\tau)} N_j^{(u)} N_j^{(\tau)} dx.$$





Final Summary

 Finite element methods solve a weak form of the exact equations in an approximate solution space.

→ < ∃→

- ► The approximate solution is defined (almost) everywhere.
- ► Neuman conditions dealt with implicitly inside formulation
- Dirichlet conditions appear in right hand side (as in finite difference methods).



References

J. Donea & A. Huerta Finite Elements Methods for Flow Problems. Wiley 2003

J.N. Reddy & D.K. Gartling The Finite Element Method in Heat Transfer and Fluid Dynamics CRC Press 1994

ㅋㅋ ㅋㅋㅋ

AMCG

Fluidity Manual AMCG 2014